



Children's Health

Measure	Age Range	Reported	Compliance Met	Exclusions
Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents – BMI (WCC)	3 years - 17 years of age	Hybrid	Record of BMI value recorded during the measurement year	
Childhood Immunization Status - Combo 10 (CIS)	2 years of age during measurement year	Hybrid	Combination of 10 vaccines completed prior to 2 nd birthday	Anaphylactic shock as a result of vaccine administered
Well-child visits in first 15 months of life (W15)	Turn 15 months during measurement year	Hybrid	6 well-care visits completed prior to 15 months of age	
Well-child visits in ages 3-6 years of life (W34)	3 years - 6 years of age	Hybrid	Yearly well-care visit	
Immunizations for Adolescents - Combo 2 (IMA)	Turn 13 years of age during measurement year	Hybrid	Combination of 3 vaccines completed prior to 13 th birthday	Anaphylactic shock as a result of vaccine administered
Adolescent Well-care visits (AWC)	12 years - 21 years of age during measurement year	Hybrid	Yearly well-care visit	

Well-care Visit Timeline:

W15 – 6 or more well-care visits before 15 months of age

W34 – yearly well-care visit for children aged 3 years to 6 years of age

AWC – yearly well-care visit for children aged 12 years to 21 years of age

Immunization Timeline:

CIS – Combo 10: series of 10 vaccines to be completed prior to 2nd birthday

- DTap, IPV, MMR, HiB, HepB, VZV, PCV, HepA, RV, Influenza

IMA – Combo 2: series of 3 vaccines to be completed prior to 13th birthday:

- Tdap – at least one administered with DOS on or between member's 10th and 13th birthday
- Meningococcal Serogroups A, C, W, Y – at least one vaccine with DOS on or between the member's 11th and 13th birthday
- HPV – at least 2 HPV vaccines with different DOS on or between the member's 9th and 13th birthday
 - There must be at least 146 days between HPV vaccines

Women's Health

Measure	Age Range	Reported	Compliance Met	Exclusions
Cervical Cancer Screening (CCS)	Women 21 years - 64 years as of December 31 of the measurement year	Hybrid	Women 21 – 64 years of age who had a Pap test 2 years prior to the measurement year or by December 31 of the measurement year. Women age 30 to 64 years of age who had a Pap test with HPV co-testing performed on the same day, every 5 years.	Evidence of a hysterectomy with no residual cervix, cervical agenesis or acquired absence of cervix.
Chlamydia Screening in Women (CHL)	Women 16 years - 24 years as of December 31 of the measurement year	Admin	Women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year. Two possible methods to identify sexually active women: pharmacy data and claim/encounter data.	Women who fell in the denominator based on a pregnancy test alone and also had: a prescription for Isotretinoin or an x-ray on the date of the pregnancy test/6 days after the pregnancy test are excluded from the measure.
Breast Cancer Screening (BCS)	Women 50 years - 74 years as of December 31 of the measurement year	Admin	Women 50-74 years of age who had a mammogram any time on or between October 1st, 2 years prior to the measurement year or by December 31 of the measurement year.	Exclusions: Women who had a bilateral mastectomy <i>Note: Biopsies, breast ultrasounds and MRIs do not count. HEDIS® does not consider them to be appropriate primary screening methods</i>
Prenatal and Postpartum Care – Prenatal (PPC)	None specified	Hybrid	Prenatal care visit in the first trimester or within 42 days of enrollment. Patient MUST be in the first trimester and Dx of pregnancy must be present in not. Documentation in the medical record must include a note indicating when the prenatal care visit occurred and evidence of the following: a basic physical OB exam, evidence that a prenatal care procedure was preformed	
Prenatal and Postpartum Care – Postpartum (PPC)	None specified	Hybrid	Deliveries of live births on or between November 6 of the year prior to the measurement year and November 5 of the measurement year are included in the measure. Postpartum care visit to an OB/GYN, midwife, FP or other PCP between 21 and 56 days after delivery with the following: pelvic exam, evaluation of weight, BP, breast and abdomen, notation of postpartum care.	



Behavioral Health

Measure	Age Range	Reported	Denominator Criteria	Compliance Met	Exclusions
Antidepressant Medication Management – Acute (AMM)	18 years of age or older as of April 30 of the measurement year	Admin	<ol style="list-style-type: none"> 12-month window starting May 1 the year prior to the measurement year ending April 30 Index prescription start date which is the earliest dispensing date for antidepressant medication Continuously enrolled for 105 days prior to index prescription start through 231 days after 	Percentage of members who remained on antidepressant medication for at least 84 days, allowable gap in medication coverage not to exceed 31 days	Members who had antidepressant medication 105 days prior to date of first prior index prescription dispense date Members without an encounter that includes a diagnosis of depression within +/- 60 days of index date
Antidepressant Medication Management – Continuation (AMM)	18 years of age or older as of April 30 of the measurement year	Admin	<ol style="list-style-type: none"> 12-month window starting May 1 the year prior to the measurement year ending April 30 Index prescription start date which is the earliest dispensing date for antidepressant medication Continuously enrolled for 105 days prior to index prescription start through 231 days after 	Percentage of members who remained on antidepressant medication for at least 180 days, allowable gap in medication coverage not to exceed 52 days	Members who had antidepressant medication 105 days prior to date of first prior index prescription dispense date Members without an encounter that includes a diagnosis of depression within +/- 60 days of index date
Follow-Up Care for Children Prescribed ADHD Medication - Initiation Phase (ADD)	6 years as of March 1 the year prior to the measurement year to 12 years as of the last day of February of the measurement year	Admin	<ol style="list-style-type: none"> 12-month window starting March 1 the year prior to the measurement year ending last day of February Prescribed ADHD medication 	Percentage of members dispensed ADHD medication who had 1 follow-up visit with practitioner w/prescribing authority during 30-day window after index date	Any member with an acute inpatient encounter for mental health of chemical dependency during the 30 days after index date
Follow-Up Care for Children Prescribed ADHD Medication - Continuation Phase (ADD)	6 years as of March 1 the year prior to the measurement year to 12 years as of the last day of February of the measurement year	Admin	<ol style="list-style-type: none"> 12-month window starting March 1 the year prior to the measurement year ending last day of February Prescribed ADHD medication 	Percentage of members dispensed ADHD medication, who remained on for at least 210 days and is compliant for the Initiation Phase had at least 2 follow-up visits with any practitioner within 270 days after the Initiation Phase ended	Any member with an acute inpatient encounter for mental health of chemical dependency during the 30 days after index date



Acute/ Chronic Care

Measure	Age Range	Reported	Compliance Met	Exclusions
Adult Body Mass Index Assessment (ABA)	18 years as of January 1 the year prior to the measurement year to 74 years as of December 31 of the measurement year	Hybrid	BMI reading taken in the measurement year or year prior	Pregnant members
Controlling High Blood Pressure (CBP)	18 years - 85 years of age as of December 31 of the measurement year	Hybrid	Blood pressure reading during the latest DOS for the measurement year <140/90	Member 66-80 years and older with advance illness and frailty
Comprehensive Diabetes Care - A1c Test (CDC-Test)	Diabetic members aged 18 years - 75 as of December 31 of the measurement year	Hybrid	A1c test completed during the measurement year	Member 66-80 years and older with advance illness and frailty
Comprehensive Diabetes Care - Poor Control (CDC-Poor)	Diabetic members aged 18 years - 75 as of December 31 of the measurement year	Hybrid	A1c test results >9% (lower score indicates better performance)	Member 66-80 years and older with advance illness and frailty
Asthma Medication Ratio (AMR)	5 years - 64 years of age as of December 31 of the measurement year	Admin	Ratio of controller inhaler use over total asthma medications, 50% or better for compliance	Advance respiratory conditions Members with no asthma medications dispensed during measurement year
Plan All-Cause Readmission (PCR)	18 years - 64 years of age as of the Index Discharge Date.	Admin	Count of unplanned readmissions within 30 days of the discharge date from hospital stay divided by total hospital discharges (lower score indicates better performance)	Member death Pregnant members